

BLS

ALS

<p>For a <u>conscious</u> patient:</p> <ul style="list-style-type: none">• Reassure, encourage coughing• O₂ prn <p>For inadequate air exchange: airway maneuvers (AHA)</p> <ul style="list-style-type: none">• Abdominal thrusts• Use chest thrusts in the obese or pregnant patient <p>NOTE: 5 Back Blows and Chest thrusts for infants <1 year. MR prn</p> <p>If patient becomes unconscious OR is found unconscious: Begin CPR</p> <p>Once obstruction is removed:</p> <ul style="list-style-type: none">• O₂ Saturation prn• High flow O₂, ventilate prn <p>NOTE: If suspected epiglottitis:</p> <ul style="list-style-type: none">• Place patient in sitting position• Do not visualize the oropharynx <p>STAT transport Treat as per Respiratory Distress Protocol S-167.</p>	<p>If patient becomes unconscious or has a decreasing <u>LOC</u>:</p> <p>Direct laryngoscopy and Magill forceps <u>SO</u>. MR prn</p> <p>Once obstruction is removed:</p> <p>Monitor EKG</p> <p>IV/IO <u>SO</u> adjust prn</p>
---	--

Note: If unable to secure airway, transport STAT while continuing CPR (unconscious patient).

Document revised: 7/1/2013
Approved:



EMS Medical Director