SVD LIBER TOO

and other STDs can harm your baby GET TESTED. GET TREATED. SPREAD THE WORD.

For confidential testing or information: www.STDSanDiego.org STD.HHSA@sdcounty.ca.gov

SYPHILIS IN WOMEN

- Syphilis in women is increasing in San Diego County.
- If a woman has syphilis while pregnant, it can lead to birth defects, miscarriage, premature birth, or stillbirth.
- Good News: Syphilis can be easily treated and cured with antibiotics, even during pregnancy.

IF YOU ARE PREGNANT:

- Prenatal care is very important to keep you and your baby healthy.
- You should be tested for syphilis three times during your pregnancy 1) at your first prenatal visit, 2) early in the 3rd trimester (around 28 weeks) and 3) at delivery.
- Call 2-1-1 San Diego or visit www.STDSanDiego.org if you don't have a regular doctor and need to find out where to get tested and treated.

Adapted from Los Angeles County Department of Public Health











SYPHILIS

Syphilis and other STDs are on the rise in San Diego County **GET TESTED. GET TREATED. SPREAD THE WORD.**

For confidential testing or information: www.STDSanDiego.org |STD.HHSA@sdcounty.ca.gov

SYPHILIS SYMPTOMS

- The first symptom is a painless sore that may appear inside or near the mouth, genitals, or anal area.
- A second symptom may be a rash that often appears on the palms of the hands, bottoms of the feet, or the skin between the armpits and upper thighs.
- Syphilis symptoms may appear and disappear.
- To cure syphilis, medical testing and treatment are required.

TEST TODAY

- If you are experiencing signs or symptoms.
- If you are getting tested for other STDs.
- If you are sexually active (testing recommended every 6-12 months).
- If you have sex without a condom (testing recommended every 6 months).
- If you have more than one sex partner (testing recommended every 6 months).
- If you are a man having sex with other men (testing recommended every 3-6 months).
- If you are pregnant or plan to get pregnant in the future.