Rotavirus



ROTAVIRUS

Rotavirus is a virus that affects the stomach and intestines. It is most common in infants and young children. Older children and adults can also get sick from rotavirus.

Since rotavirus can cause severe vomiting and diarrhea, it can lead to dehydration (loss of bodily fluids). Children who become dehydrated may need to be hospitalized.

TRANSMISSION

Rotavirus spreads easily among infants and young children. Kids can get infected when they:

- Put their fingers in their mouths after touching a contaminated surface (like a toy), or
- Eat contaminated food.



SIGNS AND SYMPTOMS

The most common symptoms of rotavirus are:

- Severe watery diarrhea,
- Vomiting,
- Fever, and
- Belly cramps.



Additional symptoms may include loss of appetite and dehydration. These symptoms can be dangerous for babies and young children.

PREVENTION

Good hygiene (like handwashing) and cleaning shared toys are important, but are not enough to control the spread of disease.

Rotavirus vaccination is the best way to protect your children, family, and community from rotavirus disease.



The rotavirus vaccine is a liquid given by mouth to babies. Depending on the vaccine brand, one dose is given at:

- · 2 and 4 months, or
- 2, 4, and 6 months.

Call your child's doctor to schedule their rotavirus vaccine today!







CALL: 1 (866) 358-2966 OPTION 5
EMAIL: PHS-IZPHN.HHSA@SDCOUNTY.CA.GOV

