



To: CAHAN San Diego Participants

Date: May 23, 2022

From: Public Health Services

Health Advisory: Human Monkeypox

Key Messages

- The Centers for Disease Control and Prevention (CDC) has issued a health alert due to a case of human monkeypox identified in Massachusetts following travel to Canada.
- Recently, monkeypox clusters have been identified in several countries with cases reporting no travel to endemic locations.
- Cases have been identified among men who have had close or intimate in-person contact with other men. The rash may be confused with other sexually transmitted infections such as secondary syphilis.

Situation

Cases of monkeypox have previously been identified in travelers from, or residents of, West African or Central African countries, where monkeypox is endemic. Since May 14, 2022, clusters of monkeypox cases have been reported in several countries that do not normally have monkeypox, including several European countries and Canada. Although previous cases outside of Africa have been associated with travel from Nigeria, most of the recent cases do not have direct travel-associated exposure risks. The United Kingdom Health Security Agency (UKHSA) was the first to announce identification of a recent U.K. case that occurred in a traveler returning from Nigeria.

Background

The Massachusetts Department of Public Health and the Centers for Disease Control and Prevention (CDC) are investigating a confirmed case of [monkeypox in the United States](#). On May 17, 2022, skin lesions that had several features suspicious for monkeypox—firm, well circumscribed, deep-seated, and umbilicated lesions—on a Massachusetts resident who recently traveled to Canada. Testing at CDC confirmed the patient was infected with a West African strain of monkeypox virus. The patient is currently isolated and does not pose a risk to the public.

On May 20, 2022, CDC issued a [Health Alert Network \(HAN\) Health Advisory](#) to ask clinicians in the United States to be vigilant to the characteristic rash associated with monkeypox. Suspicion for monkeypox should be heightened if the rash occurs in people who: 1) traveled to countries with recently confirmed cases of monkeypox; 2) reported having had contact with a person or people who have a similar appearing rash, or received a diagnosis of confirmed or suspected monkeypox; or 3) is a man who regularly has had close or intimate in-person contact with other men. Lesions may be disseminated or located on the genital or perianal area alone. Some patients may present with proctitis, and their illness could be clinically confused with a sexually transmitted infection (STI) like syphilis or herpes, or with varicella zoster virus infection. Transmission most efficiently occurs through large respiratory droplets, which requires prolonged face-to-face exposure, and direct contact with rash lesions or body fluids. Lack of a fever prodrome is not a reliable way to rule out a characteristic rash.

Actions Requested

- 1)** Consider a diagnosis of monkeypox in people who present with the characteristic rash and who in the month preceding illness:
 - a) have traveled to countries where monkeypox cases have been recently reported;
 - b) have had direct or indirect contact with someone who is arriving/returning from Africa, has a similar rash, and/or has received a diagnosis of confirmed or suspected monkeypox; or
 - c) are men who have had close or intimate in-person contact with other men.
- 2) Report suspected cases immediately by telephone** to the San Diego County Epidemiology Unit, by calling 619-692-8499 (8 am to 5 pm, Monday through Friday); 858-565-5255 (after hours) prior to testing.
 - a) Confirmatory monkeypox virus-specific testing requires a dry lesion swab specimen. [Collect](#) multiple specimens from different lesions for preliminary and confirmatory testing.

Resources

[Monkeypox | World Health Organization](#)

[CDC Health Alert: Monkeypox Virus Infection in the United States and Other Non-endemic Countries—2022](#)

[2022 United States Monkeypox Case | Monkeypox | Poxvirus | CDC](#)

[Clinical Recognition of Monkeypox | CDC](#)

[Smallpox & Other Orthopoxvirus-Associated Infections 2020 Yellow Book | CDC](#)

[Preparation and Collection of Specimens | Monkeypox | Poxvirus | CDC](#)

[Treatment | Monkeypox | Poxvirus | CDC](#)

[Infection Control Measures to Prevent Monkeypox Transmission in Hospitals | CDC](#)

Thank you for your participation.

CAHAN San Diego

County of San Diego Health & Human Services Agency

Epidemiology and Immunization Services Branch

Phone: (619) 692-8499; Fax: (858) 715-6458

Urgent Phone for pm/weekends/holidays: (858) 565-5255

E-mail: cahan@sdcounty.ca.gov

Secure Website: <https://member.everbridge.net>

Public Website: <http://www.cahansandiego.com>

This is an official
CDC Health Advisory

Distributed via the CDC Health Alert Network
May 20, 2022, 11:30 AM ET
CDCHAN-0466

**Monkeypox Virus Infection in the United States and Other
Non-endemic Countries—2022**

Summary

The Massachusetts Department of Public Health and the Centers for Disease Control and Prevention (CDC) are investigating a confirmed case of monkeypox in the United States. On May 17, 2022, skin lesions that had several features suspicious for monkeypox—firm, well circumscribed, deep-seated, and umbilicated lesions—on a Massachusetts resident prompted specialized Laboratory Response Network (LRN) testing of swab specimens collected from the resident; preliminary testing confirmed the presence of DNA consistent with an orthopoxvirus using Orthopoxvirus generic and non-variola Orthopoxvirus real-time polymerase chain reaction (PCR) assays. This group of viruses includes monkeypox virus (the causative agent of monkeypox). Testing at CDC on May 18 confirmed the patient was infected with a West African strain of monkeypox virus. The patient is currently isolated and does not pose a risk to the public.

Cases of monkeypox have previously been identified in travelers from, or residents of, West African or Central African countries where monkeypox is considered to be endemic. CDC is issuing this Health Alert Network (HAN) Health Advisory to ask clinicians in the United States to be vigilant to the characteristic rash associated with monkeypox. Suspicion for monkeypox should be heightened if the rash occurs in people who 1) traveled to countries with recently confirmed cases of monkeypox, 2) report having had contact with a person or people who have a similar appearing rash or received a diagnosis of confirmed or suspected monkeypox, or 3) is a man who regularly has close or intimate in-person contact with other men, including those met through an online website, digital application (“app”), or at a bar or party. Lesions may be disseminated or located on the genital or perianal area alone. Some patients may present with proctitis, and their illness could be clinically confused with a sexually transmitted infection (STI) like syphilis or herpes, or with varicella zoster virus infection.

Background

Since May 14, 2022, clusters of monkeypox cases, have been reported in several countries that don't normally have monkeypox. Although previous cases outside of Africa have been associated with travel from Nigeria, most of the recent cases do not have direct travel-associated exposure risks. The United Kingdom Health Security Agency (UKHSA) was the first to announce on May 7, 2022, identification of a recent U.K. case that occurred in a traveler returning from Nigeria. On May 14, 2022, UKHSA announced an unrelated cluster of monkeypox cases in two people living in the same household who have no history of recent travel. On May 16, 2022, UKHSA announced a third temporally clustered group of cases involving four people who self-identify as gay, bisexual, or men who have sex with men (MSM), none of whom have links to the three previously diagnosed patients. Some evidence suggests that cases among MSM may be epidemiologically linked; the patients in this cluster were identified at sexual health clinics. This is an evolving investigation and public health authorities hope to learn more about routes of exposure in the coming days.

Monkeypox is a zoonotic infection endemic to several Central and West African countries. The wild animal reservoir is unknown. Before May 2022, cases outside of Africa were reported either among people with recent travel to Nigeria or contact with a person with a confirmed monkeypox virus infection. However, in May 2022, nine patients were confirmed with monkeypox in England; six were among persons without a history of travel to Africa and the source of these infections is unknown.

Monkeypox disease symptoms always involve the characteristic rash, regardless of whether there is disseminated rash. Historically, the rash has been preceded by a prodrome including fever, lymphadenopathy, and often other non-specific symptoms such as malaise, headache, and muscle aches. In the most recent reported cases, prodromal symptoms may not have always occurred; some recent cases have begun with characteristic, monkeypox-like lesions in the genital and perianal region, in the absence of subjective fever and other prodromal symptoms. For this reason, cases may be confused with more commonly seen infections (e.g., syphilis, chancroid, herpes, and varicella zoster). The average incubation period for symptom onset is 5–13 days.

The typical monkeypox lesions involve the following: deep-seated and well-circumscribed lesions, often with central umbilication; and lesion progression through specific sequential stages—macules, papules, vesicles, pustules, and scabs. Synchronized progression occurs on specific anatomic sites with lesions in each stage of development for at least 1–2 days. The scabs eventually fall off¹. Lesions can occur on the palms and soles, and when generalized, the rash is very similar to that of smallpox including a centrifugal distribution. Monkeypox can occur concurrently with other rash illnesses, including varicella-zoster virus and herpes simplex virus infections. Case fatality for monkeypox is reported to range between 1 and 11%. Confirmatory laboratory diagnostic testing for monkeypox is performed using real-time polymerase chain reaction assay on lesion-derived specimens.

A person is considered infectious from the onset of symptoms and is presumed to remain infectious until lesions have crusted, those crusts have separated, and a fresh

layer of healthy skin has formed underneath. Human-to-human transmission occurs through large respiratory droplets and by direct contact with body fluids or lesion material. Respiratory droplets generally cannot travel more than a few feet, so prolonged face-to-face contact is required. Indirect contact with lesion material through fomites has also been documented. Animal-to-human transmission may occur through a bite or scratch, preparation of wild game, and direct or indirect contact with body fluids or lesion material.

There is no specific treatment for monkeypox virus infection, although antivirals developed for use in patients with smallpox may prove beneficial². Persons with direct contact (e.g., exposure to the skin, crusts, bodily fluids, or other materials) or indirect contact (e.g., presence within a six-foot radius in the absence of an N95 or filtering respirator for ≥3 hours) with a patient with monkeypox should be monitored by health departments; depending on their level of risk, some persons may be candidates for post-exposure prophylaxis with smallpox vaccine under an Investigational New Drug protocol after consultation with public health authorities.

Recommendations for Clinicians

- If clinicians identify patients with a rash that could be consistent with monkeypox, especially those with a recent travel history to a country where monkeypox has been reported, monkeypox should be considered as a possible diagnosis. The rash associated with monkeypox involves vesicles or pustules that are deep-seated, firm or hard, and well-circumscribed; the lesions may umbilicate or become confluent and progress over time to scabs. Presenting symptoms typically include fever, chills, the distinctive rash, or new lymphadenopathy; however, onset of perianal or genital lesions in the absence of subjective fever has been reported. The rash associated with monkeypox can be confused with other diseases that are more commonly encountered in clinical practice (e.g., secondary syphilis, herpes, chancroid, and varicella zoster). However, a high index of suspicion for monkeypox is warranted when evaluating people with the characteristic rash, particularly for the following groups: men who report sexual contact with other men and who present with lesions in the genital/perianal area, people reporting a significant travel history in the month before illness onset or people reporting contact with people who have a similar rash or have received a diagnosis of suspected or confirmed monkeypox.
- Information on infection prevention and control in healthcare settings is provided on the CDC website: [Infection Control: Hospital | Monkeypox | Poxvirus | CDC](#). CDC is currently reviewing this information to consider the need for updates.
- Clinicians should consult their state health department ([State Contacts](#)) if they suspect monkeypox; if the health department cannot be reached, CDC can be contacted through the CDC Emergency Operations Center (770-488-7100) as soon as monkeypox is suspected.

- All specimens should be sent through the state and territorial public health department, unless authorized to send them directly to CDC.

Recommendations for Health Departments

- If monkeypox is suspected, CDC should be consulted through the CDC Emergency Operations Center (770-488-7100).
 - Appropriately collected samples can be sent to CDC or an appropriate Laboratory Response Network laboratory for testing by PCR.
- Laboratory Response Network laboratories can provide orthopoxvirus testing on lesion specimens that clinicians obtain from suspected patients; confirmatory monkeypox virus-specific testing at CDC requires a dry lesion swab specimen. Collect multiple specimens for preliminary and confirmatory testing as follows: 1) Vigorously swab or brush lesion with two separate sterile dry polyester or Dacron swabs; 2) Break off end of applicator of each swab into a 1.5- or 2-mL screw-capped tube with O-ring or place each entire swab in a separate sterile container. Do not add or store in viral or universal transport media.
- After diagnosis of monkeypox, begin contact tracing of individuals who may have been exposed to the patient while the patient was symptomatic. Contacts should be monitored for 21 days after their last date of contact with the patient.
- Share this HAN Health Advisory with relevant healthcare provider networks, including STI clinics that may not always receive CDC HAN messages.

Recommendations for the Public

- Based on limited information available at this time, risk to the public appears low. Some people who may have symptoms of monkeypox, such as characteristic rashes or lesions, should contact their healthcare provider for a risk assessment. This includes anyone who 1) traveled to countries where monkeypox cases have been reported, 2) reports contact with a person who has a similar rash or received a diagnosis of confirmed or suspected monkeypox, or 3) is a man who has had close or intimate in-person contact with other men in the past month, including through an online website, digital application (“app”), or at a bar or party.

For More Information

- Contact your state or local health department if you have any questions or suspect a patient may have monkeypox.

- CDC Poxvirus and Rabies Branch: poxvirus@cdc.gov or for issues that cannot be resolved through emails, CDC's 24/7 Emergency Operations Center (EOC): 770-488-7100 or CDC-INFO (1-800-232-4636).

References

¹ [Clinical Recognition of Monkeypox](#)

² [Antivirals](#)

The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national, and international organizations.

Categories of Health Alert Network messages

Health Alert: Requires immediate action or attention; highest level of importance.

Health Advisory: May not require immediate action; provides important information for a specific incident or situation.

Health Update: Unlikely to require immediate action; provides updated information regarding an incident or situation.

HAN Info Service: Does not require immediate action; provides general public health information.

##This message was distributed to state and local health officers, state and local epidemiologists, state and local laboratory directors, public information officers, HAN coordinators, and clinician organizations##