



INFLUENZA WATCH

Overview

Seasonal influenza activity has been very low in San Diego. The social distancing, hand washing, and mask wearing interventions instituted to reduce the transmission of SARS-CoV-2, the virus that causes COVID-19, can prevent influenza. This will be the last issue of the Influenza Watch for the 2020-21 season.

Report Content Links

- Page 1: Current Season
- Page 2: [Communiqué](#)
[Influenza Cases by Type](#)
- Page 3: [Activity Level Indicator](#)
- Page 4: [Influenza Epi Curve](#)
- Page 5: [Influenza Cases by Age](#)
[SDIR Immunizations](#)
- Page 6: [Syndromic Indicators](#)
- Page 7: [ED ILI% and P&I Deaths](#)
- Page 8: [Deaths](#)
- Page 9: [Reporting Information](#)

Key Points

Current Week 8 (ending 2/27/2021)

- 5 new influenza detections reported
- One influenza-related death reported
- 2% influenza-like illness (ILI) among emergency department visits
- 21% of death certificates registered with pneumonia and/or influenza

2020-21 FYTD Season Summary

701


Total Cases

2

Deaths[†]

0

Outbreaks*



Virus Characteristics

- Influenza A, subtype unknown
- Influenza A (H1N1)pdm09
- Influenza A (H3)
- Influenza B, subtype unknown
- Influenza B/Victoria
- Influenza B/Yamagata
- Influenza, type unknown

[†] Flu deaths less than 18 years of age are reportable to CDPH.

* In a congregate living setting, outbreaks are defined as at least one laboratory-confirmed influenza case in the setting of a cluster (≥2 cases) of influenza-like illness (ILI) within a 72-hour period.

Table 1. Influenza Surveillance Indicators.

Indicator	2020-21 Season			2019-20 Season			Prior 3-Year Average*		
	Week 8	Week 7	Total to Date	Week 8	Total To Date	Season Total	Week 8	Total To Date	Season Total
All influenza detections reported (rapid or PCR)	5	13	701	1,212	17,909	20,711	763	13,600	17,073
Percent of emergency department visits for ILI	2%	2%		8%			7%		
Percent of deaths registered with pneumonia and/or influenza	21%	22%		9%			9%		
Number of influenza-related outbreaks [∞]	0	0	0	5	55	62	3	59	69
Number of influenza-related deaths reported [^]	1	0	2	11	74	108	10	132	176

Influenza season is July 1 – June 30, Weeks 27-26. Previous weeks case counts or percentages may change due to delayed processing or reporting.

*Includes FYs 2017-18, 2018-19, and 2019-20.

[∞]At least one case of laboratory-confirmed influenza in a setting experiencing two or more cases of influenza like illness (ILI) within a 72-hour period.

Total confirmed influenza outbreaks in prior seasons: 119 in 2017-18, 25 in 2018-19, and 62 in 2019-20.

[^]Current FY deaths are shown by week of report; by week of death for prior FYs. Total deaths reported in prior seasons: 343 in 2017-18, 77 in 2018-19, and 108 in 2019-20.

Highly Pathogenic Avian Influenza (HPAI)

Although avian influenza A viruses usually do not infect people, rare cases of human infection with these viruses have been seen. Certain strains of avian influenza A viruses can produce severe illness and death when they infect humans, and these viruses are termed highly pathogenic avian influenza (HPAI).

Infected birds shed avian influenza virus in their saliva, mucous, and feces. Person-to-person spread of avian influenza A viruses has been reported very rarely, and when it has been reported it has been limited, inefficient, and not sustained.

HPAI Asian lineage H5N1 viruses have generally been responsible for most serious human illnesses and highest mortality. So far in 2020-2021, some H5N1, but mostly H5N8, have been the HPAI strains that have infected poultry and wild birds across many countries throughout Europe and Asia. On February 20, 2021, [Russia reported the first known H5N8 avian flu infections](#) in humans in seven poultry workers.

The Centers for Disease Control and Prevention ([CDC](#)) currently recommends a neuraminidase inhibitor for treatment of human infection with avian influenza A viruses. Seasonal influenza vaccination will not prevent infection with avian influenza A viruses but can reduce the risk of co-infection with human and avian influenza A viruses.

Weekly Influenza Watch Surveillance Report

This is the final *Influenza Watch* surveillance report for this season.

If you have any comments or suggestions regarding the *Influenza Watch*, please email us at EpiDiv.HHSA@sdcountry.ca.gov.

A final influenza season summary will be available later this year, and the weekly *Influenza Watch* reports will resume again in the fall.


Thank you!

Table 2. Influenza Cases Reported, 2020-2021 Season*

Positive Test Type/Subtype	Week 8	Week 7	Total to Date	Percent to Date
Influenza A, subtype unknown	1	5	339	48.4%
Influenza A (H1N1)pdm09	0	0	0	0.0%
Influenza A (H3)	0	0	0	0.0%
Influenza B, subtype unknown	4	8	359	51.2%
Influenza B/Victoria	0	0	0	0.0%
Influenza B/Yamagata	0	0	0	0.0%
Influenza, type unknown	0	0	3	0.4%
Total	5	13	701	100.0%

*Season is July 1- June 30, Weeks 27-26.

Figure 1. Activity Level of Emergency Department Influenza-Like Illness (ILI), 2020-21 Season to Date.



	Minimal			Low		Moderate		High		
	LEVEL 1	LEVEL 2	LEVEL 3	LEVEL 4	LEVEL 5	LEVEL 6	LEVEL 7	LEVEL 8	LEVEL 9	LEVEL 10
	< mean	≥ mean and <1 standard deviation above mean	≥1 and <2 standard deviations above mean	≥2 and <3 standard deviations above mean	≥3 and <4 standard deviations above mean	≥4 and <5 standard deviations above mean	≥5 and <6 standard deviations above mean	≥6 and <7 standard deviations above mean	≥7 and <8 standard deviations above mean	≥8 standard deviations above mean

Influenza Activity Indicator:

The activity level compares the current week’s ED ILI% (emergency department influenza-like illness, percent of all visits) to the mean and number of standard deviations above of the mean of the ED ILI% in non-influenza season weeks (CDC disease weeks 27-39) from the current and prior four seasons.

There are 10 activity levels, classified as: Minimal (levels 1-3), Low (levels 4-5), Moderate (levels 6-7), and High (levels 8-10). An activity level of 1 corresponds to when the ED ILI% is below the mean; level 2 corresponds to when the ED ILI% is less than 1 standard deviation above the mean; level 3 corresponds to when the ED ILI% is more than 1 but less than 2 standard deviations above the mean, and so on, with an activity level of 10 corresponding to when the ED ILI% is at 8 or more standard deviations above the mean.

Figure 2. San Diego County Influenza Detections by Type and Week of Report, 2020-21 Season to Date (N=701).




Figure 3. Cumulative Influenza Cases Reported by CDC Week and Season.




Figure 4. Proportion of Influenza Cases by Age Group and Season.






Figure 5. Number of Influenza Vaccinations Administered* by CDC Week and Season.




* Influenza vaccinations administered and entered into the San Diego Immunization Registry (SDIR). Week 52 data are repeated for week 53 for seasons that do not include week 53.

Figure 6. Percent of Emergency Department Visits for ILI Chief Complaint by CDC Week and Season*.



* Week 52 data are repeated for week 53 for seasons that do not include week 53.

Figure 7. Percent of Emergency Department Visit Chief Complaints for Cough, Cold, Fever, or Respiratory Symptoms* by CDC Week, 2020-21 Season to Date.



* Respiratory category includes cough, cold symptoms, influenza-like illness, and other respiratory symptoms.

INFLUENZA WATCH





Figure 10. Influenza Deaths by Type and Season.





Figure 11. Percent of Influenza Deaths by Age Group, Vaccination Status, and Underlying Medical Condition, 2020-21 Season.



* known to be vaccinated

Influenza Reporting in San Diego County

Individual influenza cases are now reportable to Public Health as of October 1, 2019. Please report laboratory-positive influenza results to the County Epidemiology Unit by **FAX (858) 715-6458** using a [Confidential Morbidity Report Form](#), or an [Influenza Case Report Form](#), and/or a copy of the laboratory results. Also, please indicate if the patient died and/or is a resident of a congregate living facility (if known).

Influenza specimens may be sent to [Public Health Laboratory](#) (PHL) for confirmation and subtyping. Please contact PHL at **(619) 692-8500 before submitting** or for questions and use the current PHL Test Request Form found at https://www.sandiegocounty.gov/hhsa/programs/phs/phs_laboratory/. Contact the Epidemiology Unit by telephone **(619) 692-8499** or email (EpiDiv.HHSA@sdcounty.ca.gov) with questions about influenza data. Influenza outbreaks should be reported by telephone to **(619) 692-8499**.

Resource Links

- County of San Diego Epidemiology Unit www.sdepi.org
- County of San Diego [2019-20 Influenza Season Summary](#)
- *Influenza Watch* [Slide Deck](#) – A slide version of this report for presentations
- County of San Diego Immunization Unit (SDIZ) www.sdiz.org
- San Diego Regional Immunization Registry (SDIR) <http://www.sdiz.org/CAIR-SDIR/index.html>
- California Department of Public Health (CDPH) [Influenza Update](#)
- Centers for Disease Control and Prevention (CDC) [Influenza Surveillance](#)

Influenza Watch Data Sources

The following sources of data are used to produce this report:

- **Influenza case reports:** Medical providers and laboratories report individual cases of confirmed influenza via fax or electronic laboratory reporting (ELR) to Public Health Services Epidemiology Unit (Epidemiology).
- **Influenza deaths:** Hospital infection control professionals report influenza-related deaths. Pediatric flu deaths (under 18 years of age) are legally reportable in California; however, San Diego County requests that all influenza-related deaths be reported for surveillance purposes. Influenza-related deaths are also identified through death certificate registration. The County Office of Vital Records notifies Epidemiology when a new death is registered with influenza listed as a cause of death or underlying condition. In addition, influenza case reports are compared to death data for San Diego County, and matches are evaluated to determine if their influenza infection was related to the cause of death.
- **Percent pneumonia and influenza deaths:** The percentage of all deaths registered that had either pneumonia and/or influenza listed as a cause of death is obtained directly from the Vital Records VRIS data system on a weekly basis.
- **Influenza-like illness (ILI):** Electronic emergency department (ED) visit data is submitted to Epidemiology daily for 16 hospitals in San Diego County. The percent of ED visits for ILI or flu (based on chief complaints or diagnosis) is calculated for each week. Note: ILI is defined as fever (>100°F or 37.8°C) and cough and/or sore throat, in the absence of a known cause.
- **Influenza outbreaks:** In a congregate living setting, outbreaks are defined as at least one laboratory-confirmed influenza in the setting of a cluster (≥2 cases) of influenza-like illness (ILI) within a 72-hour period. Influenza outbreaks are reportable in California. Epidemiology identifies outbreaks when facilities call to report. Other potential outbreaks are identified when multiple cases share an address or have a residential address that matches a skilled nursing or long-term care facility.
- **Number of vaccines:** The San Diego Immunization Registry (SDIR) provides weekly updates on the number of flu vaccinations given based on the number of flu vaccinations registered by participating providers.

The purpose of the weekly *Influenza Watch* is to summarize current influenza surveillance in San Diego County.

Please note that reported weekly data are preliminary and may change due to delayed submissions and additional laboratory results.